## When to Keep Your Child Home Flowchart

if your child has	keep your child home until
<ul> <li>Chills</li> <li>Continuous Cough</li> <li>Difficulty Breathing</li> <li>Severe Sore Throat</li> <li>Loss of Appetite</li> <li>Headache</li> <li>Body Aches</li> <li>Extreme Fatigue/Tiredness</li> </ul>	<ul> <li>symptoms are improving and</li> <li>your child can actively participate in the program</li> </ul>
Fever (above 38.0°C by forehead/ear, or above 37.5°C STOP by mouth/armpit)	<ul> <li>temperature remains below 38.0°C (forehead/ear) or 37.5°C (mouth/armpit) for 24 hours without medication</li> </ul>
Nouseo     Vomiting	• 24 hours free of nausea and vomiting
Diarrhea/Irregular Bowel Movement (2 bouts)	• 24 hours after one regular bowel movement
<ul> <li>Skin Infection</li> <li>Undiagnosed Rash</li> <li>Goopy Infected Eyes</li> <li>Signs of Any Contagious Disease</li> </ul>	<ul> <li>a medical professional verbally confirms the condition is non-contagious, or</li> <li>24 hours have passed since they started prescribed antibiotics</li> </ul>
Parasite Related Condition     STOP	• a medical professional verbally confirms that they may return.
Open Sores from Contagious/Viral/ Bacterial Infection	• the sores have scabbed over
Consistent Thick Discharge from Nose that is Green/Yellow in Colour	<ul> <li>symptom free (runny opaque noses are considered symptom free)</li> </ul>
Any Signs of Reportable Communicable Disease as Defined by the BCCDC	<ul> <li>a medical professional gives written clearance that they may return and</li> <li>the sores have scabbed over</li> </ul>